



**Core subject: Medical Emergencies**

# Affairs of the heart

Cardiac emergencies in surgery and how to deal with them

Author: Wendy Berridge

CPD: 1 hour

## *Educational aims and objectives*

This article tackles procedures and protocols for cardiac emergencies in surgery. It talks about the symptoms, scientific/biological reasons and how to deal with them.

## *Anticipated outcomes*

The reader will leave with a far greater understanding of all types of cardiac emergencies and how to deal with these life-threatening occurrences in practice.

This article is published by FMC. The publisher's written consent must be obtained before any part of this article may be reproduced in any form whatsoever, including photocopies and information retrieval systems.

While every care has been taken in the preparation of this article, the publisher cannot be held responsible for the accuracy of the information printed herein, or any consequence arising from it. The views expressed herein are those of the author(s) and not necessarily the opinion of the publisher.

# Affairs of the heart

**Wendy Berridge** looks at cardiac emergencies in surgery and how to deal with them

## DEFINITIONS

**Angina:** Pain or discomfort in the chest due to reduced blood and oxygen flow to the heart. Myocardial Infarction (MI): Damage or death to an area of the heart muscle resulting from a reduced or blocked blood supply (also known as a coronary thrombosis or heart attack.)

Angina is a condition usually caused by the build-up of a cholesterol plaque on the inner lining of the coronary arteries. Cholesterol is a fatty chemical which is part of the outer lining of cells in the body, and a cholesterol plaque is a hard, thick substance caused by deposits of cholesterol on the artery wall. Over time, the build up of the plaque causes narrowing and hardening of the arteries, and this significantly reduces the blood flow to the heart muscle. When a person is resting the heart is beating at its resting rate; when any type of exertion is undertaken the heart is required to beat more strongly and at a faster rate to ensure that an adequate supply of oxygenated blood is being circulated. This is a case of 'supply and demand', and the more exertion, the more cardiac effort is required, and the more oxygen the heart will need. When the coronary arteries are narrowed, less blood can be supplied to the heart muscle, so a situation will be reached where the demand of the heart muscle for blood will exceed that which the coronary arteries can supply. The heart muscle will therefore

be deprived of oxygen and 'cramps', producing pain. Typically, an angina attack occurs with exertion, and subsides with rest. If the narrowing of an artery reaches a critical level, angina can occur at rest (unstable angina). A person with angina, especially 'unstable angina' has a high risk of suffering a heart attack in the future.

**Myocardial infarction:** The medical term for a heart attack. It is caused by a

## MYOCARDIAL INFARCTION

- Crushing, vice-like pain in the centre of the chest, possibly radiating to the left arm, neck or jaw. Can sometimes be described as indigestion. 30% of casualties experience no pain
- Breathlessness
- Sweating/pallor
- Irregular pulse, often missing beats
- Nausea
- Anxiety
- Light-headedness
- Collapse.

complete blockage of one of the coronary arteries, which results in the area of the heart supplied by that vessel becoming completely deprived of its blood supply causing the affected area to die. The usual cause of this is either for a clot (thrombus) to form on the wall of the blood vessel or for a plaque to rupture and cause a blockage. If the blockage occurs in a vessel supplying a large area of the heart then it is unlikely to be able to function normally and a cardiac arrest will result. If the area supplied is relatively small then it may be possible for the rest of the heart to compensate and normal cardiac function may be re-established.

## SIGNS AND SYMPTOMS

The symptoms of angina can be variable depending on the severity of the coronary disease, and can also be triggered by emotionally stressful situations that also increase the heart rate (supply and demand). However, on rest, the heart rate will slow down again and the symptoms should pass within approximately 5-10 minutes. Use of a glyceryl trinitrate (GTN) spray will help to relieve the symptoms. If the symptoms persist despite rest and the use of GTN spray the diagnosis should be reconsidered as it is likely that the casualty is having a myocardial infarction. The signs and symptoms of a myocardial infarction can also be variable and it is important to remember that 30% of heart attacks are 'silent' as they produce no obvious symptoms. A heart attack can happen at any time, including during rest. The pain is similar to that of angina, but is often more severe. Unlike angina, the symptoms of MI will not be relieved by the use of the GTN spray. If the symptoms last longer than 10 minutes, the casualty is probably experiencing a heart attack.

## MANAGEMENT: ANGINA

- Sit the casualty upright and allow them to lean slightly forward – it makes it easier to breathe. The dental chair in its upright position is ideal, otherwise sit them on the floor with their back to a wall
- Do not lie them flat as this will not help with breathing and will put the heart under more pressure
- High flow oxygen must be administered via the non-rebreathing mask and, if the casualty carries their own glyceryl trinitrate (GTN) spray, they should be encouraged to use it by delivering 1-2 sprays sublingually
- If they do not have their medication, use your own supply from the drug box
- The GTN spray works by causing the

## ANGINA

- Crushing, vice-like pain, usually in the central area of the chest, can radiate to the left arm, neck or jaw
- Breathlessness
- Sweating/pallor
- Regular, fast pulse
- Anxiety.



blood vessels to dilate which will increase the blood flow to the heart

- If the pain does not reduce despite the GTN, a myocardial infarction should be suspected and the paramedics called
- A person who has angina can tell you what is 'normal' for them during an attack
- If the symptoms are worse than normal

## AUTHOR



**Wendy Berridge** has worked within the dental profession for over 26 years and delivers CPR, medical emergencies and defibrillation training to dental practices within Yorkshire and Lincolnshire. All training qualifies

for verifiable CPD and meets with current UK Resuscitation Council and GDC guidelines. It takes place at each individual practice, allowing for a personal approach, and resulting in a more relaxed atmosphere.

Email: [info@berridgemedicaltraining.com](mailto:info@berridgemedicaltraining.com)  
 Website: [www.berridgemedicaltraining.com](http://www.berridgemedicaltraining.com)  
 Phone: 07771 590513

or less responsive to treatment, then immediate transfer to hospital is required

- If you are worried, it is also a good idea to make the call – even if it is just for reassurance
- Otherwise once the symptoms have improved, allow the casualty to rest before going home as the GTN will also have increased the blood flow to other organs and can result in a painful headache
- Because of this, and the fact that they have just experienced an angina attack, it is best to arrange for someone to collect them.

## MANAGEMENT: MYOCARDIAL INFARCTION

- Telephone for an ambulance immediately
- The casualty should be placed in the upright position as mentioned earlier and high flow oxygen administered via the non-rebreathing face mask
- It may also be beneficial to administer dispersible aspirin

• Make sure that the casualty is not allergic to aspirin and not already taking 'anti-coagulant' drugs (such as warfarin). If this is the case, allowing them to slowly chew a dispersible aspirin may help

- Aspirin reduces the clotting ability of the blood, and chewing the tablet instead of swallowing whole allows the drug to be absorbed quickly into the bloodstream through the buccal sulcus. Ensure you have dispersible aspirin. The ideal dose is 300mg, although any strength will do.

## OPEN AIRWAYS

If the casualty becomes unconscious, the airway must be opened and breathing checked. If they are breathing, they must be placed into recovery position as they may vomit. Continue giving high flow oxygen via the non-rebreathing mask. Monitor constantly and, if at any time the breathing stops, the rate becomes abnormal, or you are unsure, then start CPR immediately. **CPD**



# Feedback

## Quality control

An essential part of Continuing Professional Development is quality control. This gives you the chance to provide feedback to us on the quality of these articles. This is not essential, but please feel free to reply using the form below if you have any comments on the CPD offered in this publication. If there is insufficient room on this form for your comments, please attach extra sheets of paper.

Article title: \_\_\_\_\_

Author: \_\_\_\_\_

Comments on CPD offered:

---

---

---

---

---

---

---

---

Please post back to:

The Editorial Director  
FMC  
Hertford House  
Farm Close  
Shenley  
WD7 9AB